

# SM04755, a Potential Disease-Modifying Treatment for Tendinopathy, Modulates the Wnt Pathway via Inhibition of CLK2 and DYRK1A

Alyssa-Lauren O'Green, MS, Vishal Deshmukh, PhD, Timothy Seo, MS, and Yusuf Yazici, MD  
Samumed, LLC, San Diego, CA

Poster #1640

## Background


- Tendinopathy is associated with pain, inflammation, tendon degeneration, and failed healing. Despite the high prevalence of tendinopathy, its underlying pathogenesis is not fully understood<sup>1</sup>
- Wnt signaling plays an important role in tendinopathy<sup>2</sup> by modulating inflammation, tenocyte lineage specification, protease production, and tendon homeostasis<sup>3-4</sup>
- SM04755, a novel, topical, small-molecule Wnt pathway inhibitor, has previously been shown to inhibit inflammation, protect tenocytes, and increase tenocyte differentiation in nonclinical models<sup>5</sup>
- The mechanism of action of SM04755 leading to Wnt pathway inhibition, tenocyte differentiation and protection, and anti-inflammatory activity is described

## Conclusions

- SM04755 inhibited intranuclear kinases CLKs and DYRK1A, leading to Wnt pathway inhibition
- CLK and DYRK1A inhibition induced tenocyte differentiation and reduced tendon-destroying proteases in tenocytes
- SM04755 inhibited inflammatory signaling mediators and cytokine production
- SM04755, as a single agent, may potentially benefit symptoms and provide disease modification in tendinopathy
- Human tendinopathy trials are planned


## Results

**Figure 1. SM04755 was a potent inhibitor of Wnt signaling**




A. SW480 cells; n=4; Mean ± SD B. hMSCs; n=3; Mean ± SEM; \*P<0.05, \*\*P<0.01, \*\*\*P<0.001 vs. DMSO, t-test

**Figure 2. SM04755 was a potent inhibitor of CDC-like kinases (CLKs) and DYRK1A**




**Figure 3. Inhibition of CLKs and DYRK1A inhibited the Wnt pathway in rTDSCs**




n=3; Mean ± SEM; \*P<0.05, \*\*P<0.01, \*\*\*P<0.01 vs. siCtrl, t-test

**Figure 4. Inhibition of CLKs and DYRK1A induced tenocyte differentiation *in vitro***




n=3; Mean ± SEM; \*P<0.05, \*\*\*P<0.001 vs. siCtrl, t-test

**Figure 5. Inhibition of CLK1, 2, 4, and DYRK1A reduced catabolic protease expression *in vitro***



Rat tenocytes; n=3; Mean ± SEM; \*\*\*P<0.001 vs. siCtrl, one-way ANOVA

**Figure 6. SM04755 demonstrated anti-inflammatory effects *in vitro***



n=3; Mean ± SEM; \*P<0.05, \*\*P<0.01, \*\*\*P<0.001 vs. DMSO

## Methods

- Wnt pathway inhibition was assessed by a luciferase reporter assay in SW480 colon cancer cells (Fig. 1A)
- A kinome screen (318 kinases) was performed. Kinase inhibition was assessed by Thermo Fisher Z'-LYTE™ and LanthaScreen kinase assays (Fig. 2A)
- SM04755 and siRNA knockdown effects on gene expression in human mesenchymal stem cells (hMSCs) (Fig. 1B), rat tendon-derived stem cells (rTDSCs) (Fig. 3), and rat tenocytes (Figs. 4-5) were measured by qRT-PCR using TaqMan® primers. Gene expression was normalized to GAPDH

- SM04755 effects on serine/arginine-rich splicing factor (SRSF) phosphorylation in rTDSCs (Fig. 2B) and NF-κB and STAT3 phosphorylation in peripheral blood mononuclear cells (PBMCs) (Fig. 6A) were measured by Western blot
- SM04755 and siRNA knockdown effects on tenocyte marker expression in rTDSCs were assessed by immunostaining (Fig. 4)
- SM04755 effects on cytokine production in PBMCs stimulated with LPS were measured by MSD-based ELISA (Fig. 6B)

## References

- Scott A, et al. *J Orthop Sports Phys Ther.* 2015.
- Jelinsky SA, et al. *BMC Musculoskelet Disord.* 2011.
- Lui PP, et al. *Rheumatol (United Kingdom).* 2013.
- Deshmukh V, et al. *Osteoarthritis Cartil.* 2018.
- Deshmukh V, et al. *Arthritis and Rheum.* 2016.

All authors are employees, shareholders, or consultants of Samumed, LLC. Other disclosures are listed in the published abstract.

9360 Towne Centre Drive, San Diego, CA 92121  
info@samumed.com

